

Systematic $1/S$ study of the 2D Hubbard model at half-filling

Andrey V. Chubukov and Karen A. Musaelian

Department of Physics, University of Wisconsin - Madison,

1150 University Avenue, Madison, WI 53706

and

P.L. Kapitza Institute for Physical Problems

ul. Kosygina 2, Moscow, Russia 117334

(February 6, 2008)

Abstract

The 2D Hubbard model is extended by placing $2S$ orbitals at each lattice site and studied in a systematic $1/S$ expansion. The $1/S$ results for the magnetic susceptibility and the spectra of spin-wave excitations at half filling are consistent with the large S calculations for the Heisenberg antiferromagnet. The $1/S$ corrections to the fermionic spectrum lift the degeneracy along the edge of the magnetic Brillouin zone yielding minima at $(\pm\pi/2, \pm\pi/2)$. Relation to previous papers on the subject is discussed.

I. INTRODUCTION

The 2D Hubbard model and the closely related $t - J$ model have been extensively used recently in the context of high temperature superconductivity as the simplest models which capture at least some of the exciting physics of high- T_c superconductors. At half-filling, the large- U Hubbard model reduces to the nearest-neighbor Heisenberg antiferromagnet [1], which perfectly describes the properties of pure La_2CuO_4 [3–5], while at sufficiently strong doping, it describes a strongly correlated metal and possibly a superconductor [10,11].

One of the crucial issues for lightly doped antiferromagnets is the role of quantum fluctuations. In $La_{2-x}Sr_xCuO_4$, they destroy long-range order in the ground state already at a very small doping [3]. There is therefore a need for a systematic accounting of quantum fluctuations in the Hubbard model. For ordered Heisenberg antiferromagnets, a customary tool for this is the $1/S$ expansion. This expansion gives rather fast convergence and first order $1/S$ results are quite accurate even for $S = 1/2$ [6,7].

In this paper, we report the results of the large- S studies of quantum fluctuations in the Hubbard model at half-filling. We view the results as a basis for future considerations of quantum fluctuations in doped antiferromagnets.

To perform $1/S$ studies, we need to extend the Hubbard model to higher spins. A straightforward way to do this is to place $2S$ equivalent orbitals at each lattice site [14,15]. The hopping term remains the same as in the conventional Hubbard model, only the hopping integral t now scales with S . At the same time, the $Un_\uparrow n_\downarrow$ term is replaced by the Hund rule interaction term

$$H_{int} = \frac{U}{2} \sum_{n,\alpha,i} \psi_{i\alpha n}^\dagger \psi_{j\beta n}^\dagger \psi_{i\beta n} \psi_{j\alpha n} \quad (1)$$

Here $\alpha = 1, 2$ is the spin index, and i is the orbital index which runs over $i = 1, 2, \dots, 2S$. For $S = 1/2$, the summation over spin index is absent and (1) reduces to the conventional on-site repulsion term. For $S > 1/2$, the coupling term ensures (at large U) a parallel alignment of the spins of electrons at all orbitals, i.e., total spin S at each site, which forms a natural basis

for $1/S$ expansion, and also ensures an equal number of electrons at each site. Fluctuations of this number cost energy of the order of $US \gg t$. The interaction term is simply related to the total spin per site as $H_{int} = -U \sum_n (S_n^2 + (\mathcal{N}_n - 2S)^2)/4$ where $\mathcal{N}_n = \psi_{i\alpha n}^\dagger \psi_{i\alpha n}$ is the fermion density, and $\mathbf{S}_n = \sum_i \mathbf{S}_{in} = \frac{1}{2} \psi_{i\alpha n}^\dagger \boldsymbol{\sigma}_{\alpha\beta} \psi_{i\beta n}$ is the total spin on a given site. The fluctuations of the square of the total density are irrelevant at large U , and we have checked explicitly that one can equally well use either the eq. (1) or $-U \sum_n S_n^2$ for the interaction term. The former choice is however advantageous for computational purposes.

In the momentum space, the extended Hubbard Hamiltonian takes the form

$$H = \sum_k \epsilon_k a_{j\alpha k}^\dagger a_{j\alpha k} + \frac{U}{2N} \sum_{k,k',q} a_{i\alpha,k'}^\dagger a_{j\beta,-k'+q}^\dagger a_{i\beta,-k+q} a_{j\alpha,k} \quad (2)$$

Here $\epsilon_k = -2t(\cos k_x + \cos k_y)$, N is the total number of lattice sites, and the momentum summations extend over the first Brillouin zone $-\pi < k_x, k_y < \pi$.

The large- S expansion for magnetization and spin susceptibility of the Hubbard model has been previously studied in [15]. Although a complete agreement with the similar expansion in the Heisenberg model was obtained, we believe that several relevant contributions were missed, and the agreement is thus a bit fortuitous. Below we will point out the discrepancies with Ref. [15] explicitly. The semiclassical expansion results for the magnetization were also reproduced in [13] by means of a $1/z$ expansion, where z is the number of nearest neighbors.

II. MEAN-FIELD THEORY

We now briefly outline the results of mean-field ($S = \infty$) studies of the Hubbard model [8,9,12,13,16]. At the mean-field level, one introduces the condensate of $S_z(q = Q)$ where $Q = (\pi, \pi)$, and uses it to decouple the interaction term in (2). The resulting quadratic Hamiltonian then takes the form

$$H_{\text{MF}} = \sum_k \epsilon_k a_{i\alpha k}^\dagger a_{i\alpha k} - \frac{U \langle S_z \rangle}{2} \sum_k a_{i\alpha,k+Q}^\dagger \sigma_{\alpha\beta}^z a_{i\beta k}, \quad (3)$$

where the summations over momentum here and below are limited to the magnetic zone, i.e. half the first Brillouin zone. Note that $\langle S_z \rangle$ is the exact value of the condensate which includes all zero-point fluctuations. This quadratic Hamiltonian can be diagonalized by means of a Bogolubov transformation

$$\begin{aligned} c_{i\alpha k} &= u_k a_{i\alpha k} + v_k \sigma_{\alpha\beta}^z a_{i\beta, k+Q} \\ d_{i\alpha k} &= v_k a_{i\alpha k} - u_k \sigma_{\alpha\beta}^z a_{i\beta, k+Q} \end{aligned} \quad (4)$$

where c represents conduction, and d — valence electrons. The transformation explicitly reads

$$u_k = \left(\frac{1}{2} \left(1 + \frac{\epsilon_k}{E_k} \right) \right)^{1/2}; \quad v_k = \left(\frac{1}{2} \left(1 - \frac{\epsilon_k}{E_k} \right) \right)^{1/2}, \quad (5)$$

where $E_k = (\epsilon_k^2 + \Delta^2)^{1/2}$ and $\Delta = \frac{U\langle S_z \rangle}{2}$. The diagonalized Hamiltonian then takes the form

$$H = \sum_k E_k (c_{i\alpha k}^\dagger c_{i\alpha k} - d_{i\alpha k}^\dagger d_{i\alpha k}). \quad (6)$$

The value of the gap Δ is obtained from the self-consistency condition of the diagonalization procedure

$$\frac{1}{U} = \sum_k \frac{2S}{E_k}. \quad (7)$$

The bosonic spin-wave excitations appear in the theory as poles of the total transverse susceptibility

$$\chi^{+-}(q, q'; \omega) = i \int dt \langle TS_q^+(t) S_{-q'}^-(0) \rangle e^{i\omega t}. \quad (8)$$

To zeroth order in $1/S$, the total static susceptibility is given by a sum of bubble diagrams in Fig. 1:

$$\chi^{+-} = \frac{\chi_0^{+-}(q, \omega = 0)}{1 - U \chi_0^{+-}(q, \omega = 0)}, \quad (9)$$

$$\chi_0^{+-}(q, \omega) = \frac{S}{N} \sum_k \left(1 - \frac{\epsilon_k \epsilon_{k+q} - \Delta^2}{E_k E_{k+q}} \right) \frac{2}{E_k + E_{k+q}} \quad (10)$$

This simple form familiar from paramagnon theories exists, however, only for the static susceptibility. The total dynamical susceptibility is the solution of a 2×2 problem [16] because the antiferromagnetic ordering doubles the unit cell, and $\chi_0^{+-}(q, q'; \omega)$ is nonzero either when $q = q'$ or when $q = q' + Q$ (in the latter case $\chi_0^{+-}(q, q + Q, \omega) \sim \omega$). Also notice the overall factor of S in χ_0^{+-} . It comes from the summation over the orbital index in each bubble. Because of this factor, $U\chi_0^{+-}$ is $O(1)$, and this makes the total transverse susceptibility very different from the bare one.

For large Δ/t , one can expand in (10) and simplify (9) to

$$\chi^{+-}(q) = \frac{1}{2J(1 + \gamma_q)} \quad (11)$$

where $\gamma_q = (\cos k_x + \cos k_y)/2$ and $J = 4t^2/(2S)^2U$. As q approaches the ordering momentum Q , $\gamma_q \rightarrow -1$, and static susceptibility diverges as it, indeed, should because of the Golstone theorem.

III. $1/S$ EXPANSION FOR THE STATIC SUSCEPTIBILITY

We now proceed to the study of the fluctuation corrections to the static susceptibility. These corrections are related to the residual interaction between fermions. In the transverse channel there are two terms — the direct interaction of two fermions with opposite spins, and interaction between fermions of opposite spins mediated by magnetic fluctuations, see Fig.2. The effective Hamiltonian for fermion-magnon coupling was derived by D. Frenkel and one of us [13]; here we quote only the result:

$$\begin{aligned} H_{\text{tr}} = \sum_{k,q} & (c_{i\alpha k}^\dagger c_{i\beta, k+q} e_{\beta q}^\dagger \Phi_{cc}(k, q) + d_{i\alpha k}^\dagger d_{i\beta, k+q} e_{\beta q}^\dagger \Phi_{dd}(k, q) \\ & + c_{i\alpha k}^\dagger d_{i\beta, k+q} e_{\beta q}^\dagger \Phi_{cd}(k, q) + d_{i\alpha k}^\dagger c_{i\beta, k+q} e_{\beta q}^\dagger \Phi_{dc}(k, q) + \text{H.c.}) \delta_{\alpha, -\beta} \end{aligned} \quad (12)$$

Here $e_{\alpha q}$ is the boson annihilation operator with polarization α , and the vertex functions are given by

$$\Phi_{cc, dd}(k, q) = [\pm(\epsilon_k + \epsilon_{k+q})\eta_q + (\epsilon_k - \epsilon_{k+q})\bar{\eta}_q] / 2S$$

$$\Phi_{cd,dc}(k, q) = U \left[\left(1 - \frac{(\epsilon_k + \epsilon_{k+q})^2}{8\Delta^2} \right) \eta_q \mp \left(1 - \frac{(\epsilon_k - \epsilon_{k+q})^2}{8\Delta^2} \right) \bar{\eta}_q \right], \quad (13)$$

and η_q and $\bar{\eta}_q$ are given by

$$\eta_q = \frac{1}{\sqrt{2}} \left(\frac{1 - \gamma_q}{1 + \gamma_q} \right)^{1/4}, \quad \bar{\eta}_q = \frac{1}{\sqrt{2}} \left(\frac{1 + \gamma_q}{1 - \gamma_q} \right)^{1/4}. \quad (14)$$

In (13) we neglected all terms of the order $J(t/US)^m \ll J$ with $m \geq 2$ which is consistent with our intention to calculate corrections on the scale of magnetic interaction J . Further, unlike fermionic energies, the vertex functions do not have the overall factor of S . The second-order self-energy corrections to the fermionic propagators then will contain one power of S in the numerator due to the summation over the orbitals, and one power of S in the denominator which comes from $E_k \propto S$. Total corrections will therefore be $O(1)$ i.e. will have $1/S$ factor compared to the mean-field inverse propagators which scale as $O(S)$.

To calculate the spin susceptibility we will need the renormalized self-consistency condition, i.e. expression for the order parameter $\langle S_z \rangle$ in terms of $\Delta = U \langle S_z \rangle$. It should ensure the cancellation of the leading $O(1)$ term in the denominator of χ^{+-} . At $S = \infty$, this condition was given by eqn. (7). For $1/S$ corrections, however, we need the full expression for $\langle S_z \rangle$. Substituting (4) into the formula for the z component of the spin, we obtain after a simple algebra

$$\langle S_z \rangle = \frac{1}{N} \sum_{i,\alpha,k} \frac{\Delta}{E_k} (\langle d_{i\alpha k}^\dagger d_{i\alpha k} \rangle - \langle c_{i\alpha k}^\dagger c_{i\alpha k} \rangle) + \frac{\epsilon_k}{\Delta} (\langle c_{i\alpha k}^\dagger d_{i\alpha k} \rangle + \langle d_{i\alpha k}^\dagger c_{i\alpha k} \rangle) \quad (15)$$

At $S = \infty$, the only nonzero average is $\langle d_{i\alpha k}^\dagger d_{i\alpha k} \rangle = 1$, and using $\Delta = U \langle S_z \rangle$, we return to (7). However, at finite S , one has to calculate pair averages with the self-energy corrections. These corrections include terms which transform a conduction fermion into a valence one and vice versa after emitting or absorbing a spin wave. Accordingly, all pair averages, including $\langle c_{i\alpha k}^\dagger d_{i\alpha k} \rangle$ will be different from zero at finite S . The density diagrams are presented in Fig 3. A straightforward calculation using (12) and (13) yields

$$\langle S_z \rangle = \frac{2S}{N} \sum_k \frac{\Delta}{E_k} - \frac{1}{N^2} \sum_{k,p} \frac{4S^2 \Phi_{cd}^2}{(E_k + E_{k+p} + \Omega_p)^2} + \frac{4J}{U^2} (A - B), \quad (16)$$

where $\Omega_q = 4JS \sqrt{1 - \gamma_q^2}$ is the bare spin-wave frequency, and we defined

$$A = \frac{1}{N} \sum_q \frac{1 - \sqrt{1 - \gamma_q^2}}{\sqrt{1 - \gamma_q^2}}, \quad B = \frac{1}{N} \sum_q \frac{\gamma_q^2}{\sqrt{1 - \gamma_q^2}}. \quad (17)$$

Retaining only terms of the order of U we arrive at a well-known result for the Heisenberg model [6]

$$\langle S_z \rangle = S \left(1 - \frac{1}{SN} \sum_q \frac{1 - \sqrt{1 - \gamma_q^2}}{\sqrt{1 - \gamma_q^2}} \right). \quad (18)$$

We are now in a position to compute $1/S$ corrections to the static transverse susceptibility. Simple considerations show that to the first order in $1/S$, the RPA approach is still exact, but one has to include all $1/S$ corrections within a bubble and also take into account the renormalization of the relation between Δ and U . A special care should be given in selecting the diagrams as the valence-conduction bubble itself has the order of $1/U$. However this leading contribution gets cancelled in the denominator of (9), and the momentum dependence of χ^{+-} is given by the subleading terms in χ_0^{+-} which have the order of J/U^2 . The cancellation of the $O(U)$ terms should clearly survive in $1/S$ expansion, and we, therefore, have to keep all $1/S$ diagrams to the order $J/U^2 S$. Further, one should also consider the effect of including the direct longitudinal fermion-fermion interaction. Simple power counting arguments show that this does not give rise to an extra $1/S$ smallness because each time one includes an extra interaction, one also has to sum over the orbitals of intermediate fermions. We found however that in the large- U limit each inclusion of more than one triplet interaction does produce a smallness, not in $1/S$ but in $(t/US)^2$, because the frequency integration selects only those combinations of conduction and valence fermions for which the actual longitudinal interaction becomes small after being dressed by the Bogolubov coefficients. Altogether, we found 14 relevant diagrams. They are presented in Fig 4 - 7. The diagrams in Fig.4 contain no spin wave propagator, but two direct fermion-fermion interactions. The diagrams in Fig.5 contain self-energy corrections related to the exchange of spin-waves and no direct fermion-fermion interactions. The diagrams in Fig.6

contain one spin-wave propagator and one direct interaction between fermions with parallel spins. Finally, the diagrams in Fig.7 contain one spin-wave propagator and two interactions in the triplet channel. The latter diagrams were omitted in Ref [15] together with several $O(J/U^2S)$ terms in other diagrams. We found that altogether, the terms omitted in [15] cancel each other, but we do not believe that this could be anticipated in advance. Note that inclusion of self-energy corrections with the direct transverse interaction would lead to double-counting, as these corrections have already been taken into account in the mean field diagonalization.

Calculation of the diagrams is tedious but straightforward. We list the expressions in the Appendix and quote here only the final result. We found that the modified self-consistency condition ensures an exact cancellation of the denominator in (9) at $q = Q$ in accordance with the Goldstone theorem. The functional form of χ^{+-} is also unchanged compared to the mean-field result, but quantum fluctuations give rise to an overall renormalization factor, Z_χ . We have

$$\chi^{+-}(q, \omega = 0) = \frac{Z_\chi}{2J(1 + \gamma_q)}; \quad Z_\chi = 1 - \frac{B}{S} = 1 - \frac{1}{NS} \sum_q \frac{\gamma_q^2}{\sqrt{1 - \gamma_q^2}}. \quad (19)$$

As expected, this result coincides with the known $1/S$ expression for the static susceptibility of the Heisenberg antiferromagnet [6].

Finally, for completeness, we will also obtain the $1/S$ expansion result for the spin-wave velocity, c . From hydrodynamical considerations [17], we know that $c^2 = \rho_s/\chi_\perp$, where $\chi_\perp$ is the static susceptibility

$$\chi_\perp = \frac{1}{2} \chi^{+-}(q = 0, \omega = 0) = \frac{Z_\chi}{8J}, \quad (20)$$

and ρ_s is the spin stiffness defined by

$$\chi_{\text{st}}^{xx}(q \approx Q) = \frac{S_z^2}{\rho_s(q - Q)^2}, \quad (21)$$

Using the result for S_z and expanding (19) near $q = Q$, we find

$$\rho_s = JS^2 Z_\rho; \quad Z_\rho = 1 - \frac{2A}{S} + \frac{B}{S} = 1 - \frac{2}{NS} \sum_q \frac{1 - \sqrt{1 - \gamma_q^2}}{\sqrt{1 - \gamma_q^2}} + \frac{1}{NS} \sum_q \frac{\gamma_q^2}{\sqrt{1 - \gamma_q^2}} \quad (22)$$

Combining (20) and (22), we obtain

$$c^2 = 2J^2 Z_c^2; \quad Z_c^2 = \frac{Z_\rho}{Z_\chi} = 1 + \frac{2}{S}(B - A) = 1 + \frac{2}{NS} \sum_q [1 - \sqrt{1 - \gamma_q^2}] \quad (23)$$

which again coincides with the known spin-wave result.

IV. SELF-ENERGY CORRECTIONS TO THE FERMIONIC SPECTRUM

We now turn to the calculation of self-energy corrections to the single-particle excitation energy, E_k . In the mean-field approximation we had $E_k = \sqrt{\Delta^2 + \epsilon_k^2}$. At large U , it can be simplified to $E_k = \Delta + 4JS(\cos k_x + \cos k_y)^2$. These forms for E_k imply that the minimum of the fermion energy coincides with the whole edge of the magnetic Brillouin zone, $|k_x \pm k_y| = \pi$. This degeneracy in the position of the minima of E_k was responsible for the singular behavior of the Hubbard model at infinitesimally small doping [8]. We will show, however, that this degeneracy does not survive perturbative $1/S$ corrections, and the actual minima of the quasiparticle energy are located only at the particular points $(\pm\pi/2, \pm\pi/2)$ in the Brillouin zone. This is consistent with the results of other approaches [21–25].

Consider for definiteness the corrections to the energy of a valence fermion. As before, the dominant self-energy corrections at large U are those which include the exchange of transverse magnetic fluctuations. To first order in $1/S$, there are two such diagrams (Fig.8). Performing the frequency integration in these diagrams, we obtain for the Green function of a valence fermion

$$G^{-1}(k, \omega) = \omega + E_k - \frac{2S}{N} \sum_q \left(\frac{\Phi_{cd}^2}{\omega - E_{k+q} - \Omega_q + i\delta} + \frac{\Phi_{dd}^2}{\omega + E_{k+q} + \Omega_2 - i\delta} \right) \quad (24)$$

We now consider the two self-energy terms separately. The first term in (24) corresponds to a process when valence fermion transforms into a conduction one after emitting a spin wave. The matrix element for such process and the energy denominator scale as U and US , correspondingly. As a result, the first term yields the *momentum-independent* renormalizations of the gap, $\Delta \rightarrow \Delta(1 + A/S)$, and of the wavefunction residue, $Z = 1 - A/2S$. The latter is

in fact related to the renormalization of the sublattice magnetization [13] $\langle S_z \rangle = Z - 1/2$. This first self-energy term also has momentum-dependent contributions due to the k dependence in E_{k+q} , but they scale as J and, as we will see, are completely overshadowed by the momentum-dependent contributions from the second self-energy term.

The second self-energy term in (24) describes a process which includes only valence fermions. The vertex function for such process scales as t/S so intuitively one may conclude that the second term is less relevant than the first one. However, at resonance, $\omega = -E_k$, and the leading $O(\Delta)$ terms in the denominator get cancelled. The remaining terms scale as JS , so that the total self-energy correction behaves as $O(\Delta/S)$. More important, however, is that the k -dependent self-energy terms are not anymore suppressed by a factor J/U and, therefore, may substantially modify the shape of the excitation spectrum.

The general structure of the $1/S$ self-energy correction is rather involved, so below we will restrict calculations to the dispersion relation along the lines $|k_x \pm k_y| = \pi$ where the mean-field energy of valence fermions has a degenerate maximum. Simplifying eq.(24) at $\gamma_k = -1$ and $US/t \gg 1$ we obtain for the pole of the Green function

$$E_k^{1/S} = \Delta \left(1 - \frac{1}{S} + \frac{\beta(k)}{2S} \right), \quad (25)$$

where

$$\beta(k) = \frac{2}{N} \sum_q \frac{1 + \sqrt{1 - \gamma_q^2}}{2\gamma_{k+q}^2 + \sqrt{1 - \gamma_q^2}}. \quad (26)$$

It is not difficult to see that $\beta(k)$ does contain some momentum dependence, and, therefore, the excitation energy acquires dispersion along the edge of the magnetic Brillouin zone. We evaluated the integral (26) numerically and found that $\beta(k)$ has four equivalent minima at $(\pm\pi/2, \pm\pi/2)$. Near each minimum, one can expand in $\bar{k}_{x,y} = \pm\pi/2 - k_{x,y}$ and obtain the dispersion relation typical for the anisotropic 2D Fermi gas:

$$E_k^{1/S} = \bar{\Delta} + \frac{p_\perp^2}{2m_\perp} + \frac{p_\parallel^2}{2m_\parallel}, \quad (27)$$

Here $\bar{\Delta}$ differs from Δ due to momentum-independent self-energy corrections, and we introduced $p_\perp = (\bar{k}_x \pm \bar{k}_y)/\sqrt{2}$, $p_\parallel = (\bar{k}_x \mp \bar{k}_y)/\sqrt{2}$ (the upper sign is for $(\pi/2, \pi/2)$ and

$(-\pi/2, -\pi/2)$ and the lower sign is for the other two minima). One of the effective masses is finite already at the mean-field level, and we easily obtain from $E_k^{S=\infty}$:

$$m_{\perp} = \frac{1}{8JS} (1 + O(1/S)) \quad (28)$$

On the contrary, $m_{\parallel}$ is infinite at the mean-field level, and acquires a finite value only due to $1/S$ corrections. The numerical evaluation of this mass yields

$$m_{\parallel} \approx \frac{2S}{0.084\Delta}. \quad (29)$$

For the ratio of the effective masses we then obtain to the leading order in $1/S$: $m_{\parallel}/m_{\perp} \approx 188(JS/U)$. Notice the surprisingly large numerical coefficient.

Care has to be taken in applying the $1/S$ results for the masses to the physical case of $S = 1/2$. We already mentioned that in resonance, the diagram with only valence fermions has small energy denominator $O(JS)$ because the leading Δ term in E_{k+q} gets cancelled by the external frequency. In this situation, vertex and self-energy corrections to the internal fermion line are important as they contain large factors of U/J . A simple analysis similar to that in [13] shows that self-energy and vertex corrections scale as $O(U/JS)$ and $O(U/JS)^2$ respectively. Neglecting these corrections is therefore legitimate only if $U/JS \ll 1$ and, in a rigorous sense, our $1/S$ expansion for the fermionic energy is valid only if $S \gg U/J$ [18]. It has, however, been argued several times [13,19,20] that at large U , the actual (renormalized) interaction between fermions and spin-waves (which includes the wavefunction renormalization factor of fermions) retains its functional form (at least near magnon momentum $q = (\pi, \pi)$), but has the same order of magnitude as the bandwidth. The argument here is that for $U_{eff} \sim JS$, self-energy and vertex corrections do not generate any new energy scales besides JS . In this situation, the actual expansion parameter for our studies is $O(1)$, and we can expect that the lowest-order approximation we are using still yields at least qualitatively correct results.

We conclude this section by a brief comparison with other approaches. Vignale and Hedayati [21] considered the same perturbative corrections as we did, but resorted to self-consistent rather than perturbation analysis. They required that all corrections to the bare

dispersion of quasiparticles be on the scale of JS , and solved self-consistently Dyson equation for the quasiparticle energy. They found the band minimum at $(\pi/2, \pi/2)$, which agrees with our finding. We, however, did find the corrections of the order $O(\Delta/S)$ in perturbative calculations. Boninsegni and Manousakis [22] used variational Monte-Carlo to compute the hole dispersion in $t - J$ model. They also obtained band minima at $(\pm\pi/2, \pm\pi/2)$. The quantitative comparison with our results is, however, difficult because of the relatively small values of $t/J \leq 0.5$ considered in [22]. The minima at $(\pm\pi/2, \pm\pi/2)$ in the $t - J$ model were also found in the perturbative calculations in the small t/J limit [23]. Trugman [24] and Sachdev [25] performed variational calculations for the Hubbard and $t - J$ models respectively and also found the minima at $(\pm\pi/2, \pm\pi/2)$ at large t/J . We compared the bandwidth along $k_x + k_y = \pi$ and found good agreement between Sachdev's results and ours. Thus at $t/J = 2$, variational and our $1/S$ analysis yield $\Delta E = E(\pi, 0) - E(\pi/2, \pi/2) \approx 0.2t$, and $0.24t$ correspondingly.

V. CONCLUSIONS

To summarize, in this paper we considered a large S extension of the Hubbard model and calculated in $1/S$ expansion the leading quantum corrections to the static spin susceptibility, spin-wave velocity and hole dispersion at half-filling. We found that the mean-field degeneracy of the quasiparticle spectrum does not survive in $1/S$ calculations, and the actual hole dispersion has minima at $(\pm\pi/2, \pm\pi/2)$. Our results for magnetic parameters agree (as they should) with the $1/S$ calculations in the Heisenberg model. Indeed, at half-filling, the use of Hubbard model for magnetic calculations is not the easiest way to arrive at the final result. However, the method we presented here has a substantial advantage over other techniques in that it can be straightforwardly extended to doped antiferromagnets where quantum fluctuations are much more relevant than at half filling. The work along these lines is now in progress.

VI. ACKNOWLEDGEMENTS

It is our pleasure to thank D. Frenkel, R. Joynt and S. Sachdev for useful conversations. The research was supported in part by the Graduate School at the University of Wisconsin-Madison and Electric Power Research Institute.

VII. APPENDIX

Here we give the expressions for the diagrams in Fig. 4 - 7. The two diagrams on Fig. 4 are obviously irrelevant to magnetism, and, accordingly, cancel each other out giving contributions of $\pm \frac{J}{4U^2S}$. Two of the rest of the diagrams, namely those of Fig. 5a and 6a, contribute to order $1/US$, while the rest contribute to order J/U^2S . The explicit expressions for the first two diagrams are

$$\chi_{1a} = -\frac{1}{N^2} \sum_{k,p} \frac{1}{E_k} \left(1 - \frac{(\epsilon_k + \epsilon_{k+q})^2}{4\Delta^2} \right) \left[\frac{8S^2\Phi_{cd}^2}{(E_k + E_{k+p} + \Omega_p)^2} + \frac{4S^2\Phi_{cd}^2}{E_k(E_k + E_{k+p} + \Omega_p)} \right] \quad (30)$$

for the diagram 5a, and

$$\chi_{2a} = \frac{1}{N^2} \sum_{k,p} \frac{4S^2\Phi_{cd}^2}{E_k^2(E_k + E_{k+p} + \Omega_p)} - \frac{2J}{U^2S} (1 + \gamma_q) A - \frac{2J}{U^2S} B \quad (31)$$

for the diagram 6a. Here all the sums are over a half of the Brillouin zone, and constants A and B were defined in (17). Using the self-consistency condition (16), the total contribution from these diagrams can be immediately simplified to

$$\chi_{1a} + \chi_{2a} = \frac{1}{U} - \sum_k \frac{2S}{E_k} - \frac{2J}{U^2S} (2A - B) + \frac{2J}{U^2S} (1 + \gamma_q) A \quad (32)$$

Care also should be given to the calculation of the bare bubble, because the relation between Δ and U contains $1/S$ terms. From (10) we have

$$\chi_0 = \chi_0^{+-} = \sum_k \frac{2S}{E_k} - \frac{2J}{U^2} (1 + \gamma_q) \left(\frac{US}{\Delta} \right)^3 = \sum_k \frac{2S}{E_k} - \frac{2J}{U^2} (1 + \gamma_q) \left(1 + \frac{3A}{S} \right) \quad (33)$$

Assembling eqn (32) and (33), we obtain

$$\chi_0 + \chi_1 + \chi_2 = \frac{1}{U} \left(1 - \frac{2J}{U} (1 + \gamma_q) \right) - \frac{2J}{U^2 S} (2A - B) - \frac{4J}{U^2 S} (1 + \gamma_q) A \quad (34)$$

The expression for other diagrams are as follows

$$\begin{aligned} \chi_{1b} &= \frac{2J}{U^2 S} \frac{1}{N} \sum_k \sqrt{1 - \gamma_k^2}; & \chi_{1c} + \chi_{1d} &= \frac{6J}{U^2 S} (1 + \gamma_q) (A - B); & \chi_{1e} &= \frac{2J}{U^2 S} (1 + \gamma_q) A \\ \chi_{2b} &= -\frac{2J}{U^2 S} \gamma_q (A - B); & \chi_{2c} &= \frac{2J}{U^2 S} \gamma_q B; & \chi_{2d} &= \frac{2J}{U^2 S} (1 + \gamma_q) B; \\ \chi_{2e} &= -\frac{2J}{U^2 S} \frac{1}{N} \sum_k \sqrt{1 - \gamma_k^2}; & \chi_{2f} &= -\frac{2J}{U^2 S} (1 + \gamma_q) A. \end{aligned} \quad (35)$$

Finally, the diagrams in Fig 7 are

$$\chi_{3a} = \frac{2J}{U^2 S} A; \quad \chi_{3b} = -\frac{2J}{U^2 S} \gamma_q B. \quad (36)$$

Summing up all contributions and substituting them into eqn. (9), we obtain the result quoted in eqn. (19).

REFERENCES

- [1] J. E. Hirsch, Phys. Rev. Lett. **54**, 1317 (1985) ; C. Gros, R. Joynt, and T. M. Rice, Phys. Rev. B **36**, 381 (1987).
- [2] J. Oitmaa and D. D. Betts, Can. J. Phys. **56**, 897 (1971); D.A.Huse, Phys. Rev. B **37**, 2380 (1988); E. Manousakis, Rev. Mod. Phys. **63**, 1 (1991).
- [3] B. Keimer *et al*, Phys. Rev. B **46**, 14034 (1992).
- [4] S. Chakravarty, B. I. Halperin, and D. R. Nelson, Phys. Rev. B **39**, 2344 (1989).
- [5] A. Chubukov and S. Sachdev, Phys. Rev. Lett. **71**, 169 (1993).
- [6] T.Oguchi, Phys. Rev. **117**, 117(1960); J.Igarashi, Phys. Rev. B **46**, 10763 (1992)
- [7] J.D.Reger and A.P.Young, Phys. Rev. B **37**, 5978 (1988); R.R.P.Singh, Phys. Rev. B **39**, 9760 (1989); R.R.P.Singh and D.Huse, *ibid.* **40**, 7247 (1989).
- [8] A. Singh, Z. Tešanović, and Ju. H. Kim, Phys. Rev. B **44**, 7757 (1991).
- [9] S. John and P. Voruganti, Phys. Rev. B **43**, 10815 (1991); S. John and P. Voruganti, and W.Goff, Phys. Rev. B **43**, 13365 (1991).
- [10] P. Montoux, A. V. Balatsky, and D. Pines, Phys. Rev. Lett. **67**, 3448 (1991).
- [11] See e.g. D. J. Scalapino, E. Loh, Jr., and J. E. Hirsch, Phys. Rev. B **35**, 6694 (1987).
- [12] H.Monien and K.S.Bedell, Phys. Rev. B **45**, 3164 (1992).
- [13] A.V.Chubukov and D.M.Frenkel, Phys. Rev. B **46**, 11884 (1992).
- [14] I.Affleck and F.D.M.Haldane, Phys. Rev. B **36**, 5291 (1987).
- [15] A.Singh, Phys. Rev. B **43**, 3617 (1991).
- [16] J.R.Schrieffer, X.G.Wen, and S.C.Zhang, Phys. Rev. B **39**, 11663 (1989).
- [17] B.I.Halperin and P.C.Hohenberg, Phys. Rev. **188**, 898 (1969); D.Forster, *Hydrodynamic*

Fluctuations, Broken Symmetry, and Correlation Functions (Benjamin/Cummings, Reading, MA, 1975)

- [18] For the calculations of the static susceptibility, vertex corrections of the type discussed in SecIV are irrelevant because the external frequency is zero (or, more generally, it is $O(JS)$ for all dynamical magnetic properties).
- [19] B. I. Shraiman and E. D. Siggia, Phys. Rev. Lett. **60**, 740 (1988); **61**, 467 (1988).
- [20] C. L. Kane, P. A. Lee, and N. Read, Phys. Rev. B **39**, 6880 (1989).
- [21] G. Vignale and M.R. Hedayati, Phys. Rev. B **42**, 786 (1990).
- [22] M. Boninsegni and E. Manousakis, Phys. Rev. B **43**, 10353 (1991).
- [23] . E. Dagotto, R. Joynt, A. Moreo, S. Bacci and E. Gagliano, Phys. Rev B **41**, 9049 (1990).
- [24] S. Trugman, Phys. Rev B **37**, 1597 (1988).
- [25] S. Sachdev, Phys. Rev B **39**, 12232 (1989).

FIGURES

FIG. 1. The RPA series for the total transverse susceptibility for $S = \text{infy}$. The first term represents the simple bubble, which is the building block of the ladder.

FIG. 2. The vertex function for the exchange of magnetic fluctuations. This interaction is obtained from the direct transverse fermion-fermion coupling after a summation of ladder diagrams.

FIG. 3. The $1/S$ diagrams for the staggered magnetization. The solid line corresponds to a valence fermion, dashed line corresponds to the conduction fermion, and the wavy line represents the bosonic spin-wave propagator.

FIG. 4. The diagram with two direct fermion-fermion interaction lines in both the transverse and longitudinal channels.

FIG. 5. Diagrams for the irreducible susceptibility which do not contain a fermion-fermion interaction in the longitudinal channel. A half of the diagrams is presented. The rest are equivalent to those presented on Figs.5-7 and can be obtained by replacing simultaneously all conduction electron propagators by valence ones and vice versa.

FIG. 6. Diagrams for the irreducible susceptibility which contain one direct fermion-fermion interaction line.

FIG. 7. The same as in Fig.6 but with two fermion-fermion interaction lines in the longitudinal channel.

FIG. 8. Self-energy diagrams which contribute to the propagator of valence fermions to order $1/S$.

TABLES

TABLE I. Quasiparticle dispersion along $k_x + k_y = \pi$. Energies are in units of Δ .

$(\pi/2, \pi/2)$	$(5\pi/8, 3\pi/8)$	$(3\pi/4, \pi/4)$	$(7\pi/8, \pi/8)$	$(\pi, 0)$
0.739	0.751	0.784	0.824	0.842